

PRELIMINARY CHECK-LIST OF RED, BROWN and GREEN ALGAE FROM THE
AEGEAN SEA
based on published and unpublished records
(unconfirmed records are queried)

RHODOPHYTA Wettstein
(RHODOPHYCEAE Rabenhorst)
BANGIOPHYCIDAE Engler

PORPHYRIDIALES Skuja
PORPHYRIDIACEAE Skuja

CHROODACTYLON Hansgirg

C. ornatum (C. Agardh) Basson

STYLNEMA Reinsch

S. alsidii (Zanardini) Drew

S. cornu-cervi Reinsch

ERYTHROPELTALES Garbary, Hansen & Scagel
ERYTHROTRICHIACEAE Smith

ERYTHROCLADIA Rosenvinge

E. irregularis Rosenvinge

ERYTHROTRICHIA Areschoug

E. carnea (Dillwyn) J. Agardh

E. investians (Zanardini) Bornet

E. welwitschii (Ruprecht) Batters

? POPHYROSTROMIUM Trevisan

? P. ciliare (Carmichael) Wynne

NOTE: Reported from Saronikos Gulf (Tsiamis & al. 2010, f.30-32) with a 60 µm base,
similar to Berthold's illustration of *Erythrotrichia ciliaris* from Neapoli (= *E. bertholdii*)

SAHLINGIA Kornmann

S. subintegra (Rosenvinge) Kornmann

BANGIALES Nägeli
BANGIACEAE Engler

BANGIA Lyngbye

B. fuscopurpurea (Dillwyn) Lyngbye

NOTE: B. atropurpurea (Roth) C. Agardh apparently pertains to fresh water populations

PORPHYRACEAE Kützing

PORPHYRA C. Agardh

P. leucosticta Thuret

? P. linearis Greville

NOTE: Herbarium specimens were examined by Varela-Alvarez (pers.comm.), who questioned the identification.

PYROPIA J. Agardh

P. elongata (Kylin) Neefus & Brodie in Sutherland &al. (2011)

[*Porphyra rosengurtii* Coll & Cox (Brodie &al. 2007)]

NOTE: *Porphyra rosengurtii* was reported from Kavala (Brodie &al. 2007) and later considered conspecific with *Porphyra elongata* Kylin from the Swedish west coast (see Sutherland &al. in J.Phycol.47: 1143. 2011).

P. koreana (Hwang & Lee) Hwang &al. in Sutherland (2011) (Vergés &al. in Bot.mar.56: 229-240. 2013)

[*P. olivii* Orfanidis, Neefus & Bray in Brodie &al. (2007); type locality: Thessaloniki]

NOTE: Also known from Tergesti and localities in the NW Atlantic; it represents all (?) previous records of *P. leucosticta*

RHODOCHAETALES Skuja

RHODOCHAETACEAE Schmitz

RHODOCHAETE Thuret

R. parvula Thuret

FLORIDEOPHYCIDAE Schmitz
HILDENBRANDIALES Pueschel & Cole
HILDENBRANDIACEAE Rabenhorst

HILDENBRANDIA Nardo

H. rubra (Sommerfelt) Meneghini

CORALLINALES Silva & Johansen
CORALLINACEAE Lamouroux
CORALLINOIDAE
CORALLINEAE

CORALLINA Linné

? C. caespitosa Walker, Brodie & Irvine (in Eur.J.Phycol.45:291.2009)

NOTE: 'Nea Karvali, Greece' cited on fig.2, p. 291

C. officinalis Linné

C. elongata Ellis & Solander

JANIA Lamouroux

J. adhaerens Lamouroux

J. corniculata (Linné) Lamouroux

J. longifurca Zanardini

J. rubens (Linné) Lamouroux

HALIPTILON (Decaisne) Lindley

H. virgatum (Zanardini) Garbary & Johansen
(Corallina granifera Ellis & Solander)

LITHOPHYLLOIDEAE Setchell
(AMPHIROIDEAE Johansen)
LITHOPHYLLEAE Zanardini

AMPHIROA Lamouroux

? A. beauvoisii Lamouroux

NOTE: Reported from Smyrna (Babbini & Bressan 1997)

A. rigida Lamouroux

A. rubra (Philippi) Woelkerling
(A. cryptarthrodia Zanardini)

GONIOLITHON Foslie

G. byssoides (Lamarck) Foslie
(Nullipora trochanter Bory)

NOTE: The basionym is Nullipora byssoides Lamarck ('Système des Animaux sans Vertèbres'. 374. 1801). Lamarck described the species as '**Nullipora byssoides*. n. *Gleba corallina*. *Seba Mus.3*, t.116, f.7.'. While *Gleba corallina*...is a polynomial, Seba's ('Locupletissimi rerum naturalium thesauri...1758', pl.116, f.7) illustration was explicitly based on Mediterranean material (see text of fig. 7) and is sufficient to stand as a description because it shows typical features of this species, aiding identification (Art. 38.7-10; Melbourne Code). Such features are the compact thallus formed by erect, densely branched roundish (and not lamellate) protuberances. This illustration was included in most later accounts of the species (see e.g. Areschoug in Agardh Sp. Alg. II: 522. 1852), and is the unequivocal lectotype of Goniolithon byssoides.

G. papillosum (Hauck) Foslie

LITHOPHYLLUM Philippi

L. dentatum (Kützing) Foslie
[? L. agariciforme (Pallas) Falkenberg]

L. lichenoides (Philippi) Foslie
[L. tortuosum (Esper) Foslie sensu Lemoine]

L. incrassans Philippi

L. racemus (Lamarck) Foslie

L. stictaeforme (Areschoug) Hauck (Athanasiadis 1999b)
(L. expansum sensu Lemoine)

? L. subtenellum (Foslie) Foslie

NOTE: The type is apparently representative of L. orbiculatum (Foslie) Foslie, while older Mediterranean records can be based on Hydrolithon samoense (Foslie) Keats & Chamberlain (Irvine & Chamberlain 1994:127)

TENAREA Bory

T. tortuosa (Esper) Lemoine (Athanasiadis 1995)

TITANODERMA Nägeli

T. corallinae (Crouan frat.) Woelkerling, Chamberlain & Silva

T. pustulatum (Lamouroux) Nägeli

T. cystoseirae (Hauck) Woelkerling, Chamberlain & Silva (Athanasiadis 1989)

T. verrucatum (Lamouroux) Chamberlain

MASTOPHOROIDEAE Setchell
MASTOPHOREAE Svedelius

FOSLIELLA Howe

F. farinosa (Lamouroux) Howe

F. farinosa f. *callithamnioides* (Foslie) Chamberlain

F. farinosa var. *chalicodictya* Taylor (Sartoni & De Biasi 1999, as *Hydrolithon*)

LITHOPORELLA Foslie

L. sp.

NOTE: Two Mediterranean records are known: one from the Gulf of Taranto ('1960...dans un dragage a 60 m') and the other from Egina Island in the South Aegean Sea, collected during the 'croisiere du "Thor" 1908-1910, draguee, prof. 55 m, coll. Paulsen; echantillon unique sacrifice pour l'etude...' Lemoine (1973:44)

NEOGONIOLITHON Setchell & Mason

N. notarisii (Dufour) Hamel & Lemoine (Athanasiadis 1997a)
[*Spongitis notarisii* (Dufour) Athanasiadis (1987)]

PNEOPHYLLUM Kützing

P. confervicola (Kützing) Chamberlain

P. fragile Kützing

P. caulerpae (Foslie) Jones & Woelkerling

P. zonale (Crouan frat.) Chamberlain

SPONGITIS Kützing
(*Spongites* orth.var.)

NOTE: After the feminine substantive σπογγίτις (of stone or herb; see Liddell & Scott 'Greek-English Lexicon':1628. 1968) - not the masculine substantive σπογγίτης [sponge-like (Liddell & Scott l.c.)]. The first etymology is certain because it was later explained by Kützing (in *Phycologia germanica*: 296. 1845, 'Steintang'). Kützing also adopted consistently female gender for all species of this genus throughout his publications (see *Phycologia generalis*, *Phycologia germanica*, *Species Algarum*, etc). Hence the orthography of the generic name should be corrected to *Spongitis* (Art. 60.1, 60.2) to avoid confusion with the masculine substantive *Spongites* that is used as an epithet of species names in zoology (e.g. *Cellepora*

spongites Linné, Eschara spongites Pallas, and Millepora spongites Linné

S. ramulosa (Philippi) Kützing
(? *Neogoniolithon mamillosum* Auctorum)

HAPALIDIACEAE J.E.Gray
CHOREONEMATOIDEAE Woelkerling
CHOREONEMATEAE Svedelius

CHOREONEMA Schmitz

C. thuretii (Bornet) Schmitz

MELOBESIOIDEAE Bizzozero
MELOBESIEAE Areschoug

LITHOTHAMNION Heydrich

L. coralloides (Crouan frat.) Crouan frat.
[*Lithothamnion (Lithophyllum) solutum*]

L. crispatum Hauck (Basso &al. 2011, Asia Minor)

NOTE: Previous records (as *Lithothamnion* or *Lithophyllum*) from Tenedos, Thasos, Crete are uncertain (Athanasiadis 1987). Basso &al. (l.c.) observed a single embedded multiporate conceptacle in the lectotype having distinctively sunken rosette cells, and they proposed conspecificity with *L. superpositum* Foslie (1900, type locality: Cape of Good Hope). More characters need to be compared in these species taking also into consideration the Caribbean *Lithothamnion ruptile* Foslie (Ballantine &al. in Bot.mar.54: 295.2011).

? *L. minervae* Basso (1995)
(*Lithothamnion fruticulosum* sensu Hamel & Lemoine)

? *L. sonderi* Hauck

MELOBESIA Lamouroux

M. membranacea (Esper) Lamouroux

MESOPHYLLUM Lemoine

M. expansum (Philippi) Cabioch & Mendoza (Athanasiadis & Neto 2010)

? *M. lichenoides* (Ellis) Lemoine

M. macedonis Athanasiadis (1999a)

? *M. philippii* (Foslie) Adey
(*Lithothamnion philippii* Foslie)

PHYMATOLITHEAE Adey & Johansen

PHYMATOLITHON Foslie

P. calcareum (Pallas) Adey & McKibbin

P. lenormandii (Areschoug) Adey

ACROCHAETIALES J. Feldmann
ACROCHAETIACEAE Taylor

ACROCHAETIUM Nägeli

Acrochaetium crassipes (Børgesen) Børgesen
[*Audouinella crassipes* (Børgesen) Garbary]

A. mediterraneum (Levring) Athanasiadis (2003:106)
[*Audouinella mediterranea* (Levring) Ballesteros]

Acrochaetium molinieri Coppejans & Boudouresque
[*Audouinella molinieri* (Coppejans & Boudouresque) Garbary]

A. moniliforme (Rosenvinge) Børgesen
[*Audouinella humile* (Rosenvinge) Garbary]

A. secundatum (Lyngbye) Nägeli

COLACONEMA Batters

C. daviesii (Dillwyn) Stegenga
[*Audouinella daviesii* (Dillwyn) Woelkerling]

RHODOCHORTON Nägeli

R. purpureum (Lightfoot) Rosenvinge

NEMALIALES Schmitz
GALAXAURACEAE Parkinson

SCINAIA Bivona-Bernardi

S. furcellata (Turner) J. Agardh

TRICLEOCARPA Huisman & Borowitzka (1990)

T. fragilis (Linné) Huisman & Townsend (1993)
[*Galaxaura oblongata* (Ellis & Solander) Lamouroux]

LIAGORACEAE Kützing

GANONEMA Fan & Wang

G. farinosum (Lamouroux) Fan & Wang
(*Cladostephus dubium* Bory)

NOTE: Collected near Georgiopolis, North Crete, in the upper sublittoral (Stockholm's Hoglandsskola, klass N3, excursion in Sept. 2003; material in GB). Hamel (1931) reported that *C. dubium* Bory (type locality:Methana) was representative of this species and also found a second specimen at PC collected at Rhodes in July 1894 by Nemetz and determined as *Liagora ceranoides* by Reinbold

LIAGORA Lamouroux

L. distenta (Mertens) Lamouroux

L. viscosa (Forsskål) C. Agardh

NEMALION Duby

N. helminthoides (Velley) Batters

CERAMIALES Naegel
CERAMIINEAE ined.
DELESSERIOPSIDEAE Itono & Tanaka

BALLIELLA Itono & Tanaka

B. cladoderma (Zanardini) Athanasiadis (1987)

CERAMIACEAE Dumortier
CALLITHAMNIOIDEAE Itono
PTILOTEAE Cramer

GYMNOTHAMNION J. Agardh

G. elegans (C. Agardh) J. Agardh

SPERMOTHAMNIEAE Schmitz & Hauptfleisch

LEJOLISIA Bornet

L. mediterranea Bornet

PTILOTHAMNION Thuret

P. pluma (Dillwyn) Thuret

SPERMOTHAMNION Areschoug

S. flabellatum Bornet

S. repens (Dillwyn) Rosenvinge

COMPSOTHAMNIEAE Schmitz & Hauptfleisch

COMPSOTHAMNION (Nägeli) Schmitz

C. thyoides (Smith & Sowerby) Nägeli

NOTE: The originally spelling was *thuioides* - after θυια or θυα, the odorous cedar Juniperus foetidissima (Theophrastus, H.P.1.9.3, 4.1 - Liddell & Scott in Greek-English Lexicon: 808. 1985). It was later corrected to *thujoides* by S.F. Gray (Nat. Arr. Br. Plants. 323. 1821), after the genus *Thuja* Linné. Agardh (Syn. Alg. Scand. xxvii. 1817) changed the spelling to *thuyoides* - possibly adopting French orthography. The latinization of the above two words is *thyia* or *thya* (Lewis & Short A Latin Dict.: 1869. 1984) and hence the correct orthography should be *thyoides* (or *thyioides*)

PLEONOSPORIUM Nägeli

P. borreri (Smith & Sowerby) Nägeli

CALLITHAMNIEAE Schmitz & Hauptfleisch

AGLAOTHAMNION Feldmann-Mazoyer

A. cordatum (Børgesen) Feldmann-Mazoyer
(*Callithamnion cordatum* Børgesen)

A. hookeri (Dillwyn) Maggs & Hommersand
[*Callithamnion hookeri* (Dillwyn) S. F. Gray]

A. tenuissimum (Bonnemaison) Feldmann-Mazoyer
(*Callithamnion byssoides* Harvey)

A. tripinnatum (C. Agardh) Feldmann-Mazoyer
(*Callithamnion decompositum* sensu Athanasiadis 1987)

CALLITHAMNION Lyngbye

C. corymbosum (Smith & Sowerby) Lyngbye

? *C. tetragonum* (Withering) S. F. Gray

? *C. granulatum* (Ducluzeau) C. Agardh

SEIROSPORA Harvey

S. interrupta (Smith & Sowerby) Schmitz

S. sp.

CERAMIOIDEAE De Toni
PTEROHAMNIEAE Athanasiadis

PTEROHAMNION Nägeli

P. crispum (Ducluzeau) Nägeli

DOHRNIELEAE Feldmann-Mazoyer

ANTITHAMNIONELLA Lyle

A. elegans (Berthold) Price & John

NOTE: Recorded from Saronikos & Thermaikos Gulfs, and probably representing a recent introduction (Tsiamis &al. 2011)

CALLITHAMNIELLA Feldmann-Mazoyer

C. tingitana (Bornet) Feldmann-Mazoyer

ANTITHAMNIEAE Hommersand

ANTITHAMNION Nägeli

A. cruciatum (C. Agardh) Nägeli

A. decipiens (J. Agardh) Athanasiadis (1996)
(*A. ogdeniae* Abbott)

A. heterocladum Funk

A. tenuissimum (Hauck) Schiffner

CERAMIEAE Duby

CENTROCERAS Kützing

C. gasparrinii (Menghini) Kützing
 NOTE: Saronikos Gulf (Tsiamis &al. 2010)

C. sp.

[*C. clavulatum* (C. Agardh) Montagne sensu lato]

NOTE: Won &al. (in J. Phycol. 45:227-250. 2009) reported *C. clavulatum* to be restricted to the Southern Hemisphere and that at least 2 other species of *Centroceras* (*C. gasparrinii* (Meneghini) Kützing & *C. micracanthum* Kützing) occur in the Mediterranean

CERAMIUM Roth

C. bertholdii Funk

C. bisporum Ballantine

(*C. codii* sensu Athanasiadis 1987, p.p.)

NOTE: Observed on *Peyssonnelia rubra* collected in 1980 at Achlada (Sithonia) and included within the wide concept of *Ceramium codii* (Athanasiadis 1987). This is the first record of the species that was originally described in 1990 (based on Caribbean collections made in 1987) and later reported in the western Mediterranean (see Sartoni & Boddi 2002)

C. ciliatum (Ellis) Ducluzeau

C. circinatum (Kützing) J. Agardh

C. codii (Richards) Mazoyer

? *C. diaphanum* (Lightfoot) Roth

C. echionotum J. Agardh

? *C. elegans* (Roth) Ducluzeau

C. flaccidum (Kützing) Ardisson

C. graecum Lazaridou & Boudouresque (1992)

NOTE: Also observed in the harbour of Porto-Kufo, in the North Aegean, suggesting that the species is not indigenous in the Mediterranean Sea (unpubl. data, material in GB)

? *C. rubrum* Auctorum

? *C. tenuicorne* (Kützing) Waern

(*C. strictum* sensu Harvey)

? *C. tenuissimum* (Roth) Areschoug

CORALLOPHILA Weber-van Bosse

C. cinnabarina (Bory) R. Norris

[*Ceramium cinnabarinum* (J. Agardh) Hauck, *Centroceras pignattii* Giaccone]

SPYRIDIEAE Schmitz & Hauptfleisch

SPYRIDIA Harvey

S. filamentosa (Wulfen) Harvey

CROUANIEAE Schmitz & Hauptfleisch

CROUANIA J. Agardh

C. attenuata (C. Agardh) J. Agardh

C. francescoi Cormaci, Furnari & Scammarca
(*C. francisci*, orth. variant)

GULSONIA Harvey

G. nodulosa (Ercegovic) J. Feldmann & G. Feldmann

? *PSEUDOCROUANIA* Funk

? *P. ischiana* Funk

WRANGELIEAE Schmitz & Hauptfleisch

WRANGELIA C. Agardh

W. penicillata (C. Agardh) C. Agardh

GRIFFITHSIEAE Schmitz & Hauptfleisch

ANOTRICHIUM Nägeli

A. barbatum (C. Agardh) Nägeli

A. furcellatum (J. Agardh) Baldock

A. tenue (C. Agardh) Nägeli

GRIFFITHSIA C. Agardh

G. opuntioides J. Agardh

G. phyllamphora J. Agardh

MONOSPORUS Solier

M. pedicellatus (Smith & Sowerby) Solier

VICKERSIA Karsakoff

V. baccata (J. Agardh) Karsakoff

NOTE: Saronikos Gulf (Tsiamis & al. 2011)

SPHONDYLOTHAMNIEAE Feldmann-Mazoyer

SPHONDYLOTHAMNION Nägeli

S. multifidum (Hudson) Nägeli

DASYACEAE Kützing

DASYA C. Agardh

D. baillouviana (Gmelin) Montagne

D. corymbifera J. Agardh

D. hutchinsiae Harvey

D. ocellata (Grateloup) Harvey

D. punicea Meneghini

EYPOGODON Kützing (Silva & al. 1987)
[*Dasyopsis* (Montagne) Montagne]

E. planus (C. Agardh) Kützing
[*D. plana* (C. Agardh) Zanardini]

E. spinellus (C. Agardh) Kützing
[*D. spinella* (C. Agardh) Zanardini]

HALYDICTYON Zanardini

(*Halodictyon*, *Halidictyon*, orth. variants)

NOTE: The original orthography *Halydictyon* was corrected to *Halodictyon* by Montagne, *Halidictyon* by Nägeli and *Halodictyon* by De Toni. Since, compound words based on the Greek $\alpha\lambda\varsigma$ (genitive $\alpha\lambda-\o\varsigma$, of the sea) may be spelled $\alpha\lambda\o$, $\alpha\lambda\iota$, or $\alpha\lambda\nu$, the original spelling should be maintained. Compare to *Halichrysis* Schmitz, *Haloplegma* Montagne, *Halopitys*, *Halosiphon*, *Halymenia* and *Halidrys*

H. mirabile Zanardini

HETEROSIPHONIA Montagne

H. crispella (C. Agardh) Wynne
DELESSERIACEAE Bory

ACROSORIUM Kützing

A. venulosum (Zanardini) Kylin

APOGLOSSUM J. Agardh

A. ruscifolium (Turner) J. Agardh

DRACHIELLA Ernst & J. Feldmann

D. minuta (Kylin) Maggs & Hommersand (1993)
(*Myriogramme minuta* Kylin)

ERYTHROGLOSSUM J. Agardh

E. sandrianum (Kützing) Kylin

HYPOGLOSSUM Kützing

H. hypoglossoides (Stackhouse) Collins & Hervey

MYRIOGRAMME Kylin

M. distromatica Boudouresque

NITOPHYLLUM Greville

? *N. confervaceum* Meneghini

N. punctatum (Stackhouse) Greville

RADICILINGUA Papenfuss

R. reptans (Kylin) Papenfuss

R. thysanorhizans (Holmes) Papenfuss

NOTE: Saronikos Gulf (Tsiamis &al. 2010)

SARCOMENIACEAE Womersley
SARCOMENIEAE J. Agardh

COTTONIELLA Børgesen

C. filamentosa var. algeriensis (Schotter) Cormaci & Furnari

NOTE: Rhodes (Tsiamis &al. 2011)

RHODOMELACEAE Areschoug
AMANSIEAE Schmitz

HALOPITYS Kützing

NOTE: After $\alpha\lambda\varsigma$ (genitive $\alpha\lambda-\o\varsigma$, of the sea) and the feminine substantive $\pi\tau\upsilon\varsigma$ (pine); ‘Seekiefer’ (Kützing in Phycologia germanica: 331. 1845). Hence the correct orthography is *Halopitys* (De Toni Syl.Alg.IV.sect.III: 1080. 1903; Preda Fl.Italica Crypt.: 190. 1909)

H. incurva (Hudson) Batters

(H. incurvus, orth. variant)

OSMUNDARIA Lamouroux

O. volubilis (Linné) R. Norris (1991)

(Vidalia volubilis (Linné) J. Agardh)

RYTIPHLOEA C. Agardh

(Rytiphlaea, orth. variant)

NOTE: The second compound after the masculine word $\phi\lambda\o\iota\o\varsigma$ (cortex), apparently latinized with female gender. Hence the correct orthography should be *Rytiphloea* (Kützing Phycologia germanica: 334. 185; De Toni Syl.Alg.IV.sect.III: 1093. 1903).

R. tinctoria (Clemente) C. Agardh

BRONGNIARTELLEAE Parsons

BRONGNIARTELLA Bory

B. byssoides (Goodenough & Woodward) Schmitz

CHONDRIEAE J. Agardh

ACANTHOPHORA Lamouroux

A. nayadiformis (Delile) Papenfuss

CHONDRIA C. Agardh

C. capillaris (Hudson) Wynne (1991)

[C. tenuissima (Withering) C. Agardh]

C. coerulescens (J. Agardh) Falkenberg

? *C. collinsiana* Howe

C. dasiphylla (Woodward) C. Agardh

C. mairei Feldmann-Mazoyer (Aysel 1983)

? *C. polyrhiza* Collins & Hervey

LAURENCIEAE Schmitz

PALISADA Nam

Palisada perforata (Bory ex Montagne) Nam (2007)

[*C. papillosum* (C. Agardh) Garbary & Harper, *Laurencia papillosa* (C. Agardh) Greville]

Palisada tenerrima (Cremades) Serio & al. (2010:14)

[*Chondrophycus tenerrimus* (Cremades) Furnari, Boisset, Cormaci & Serio (2002)]

Palisada thyoides (Kützing) Cassano, Senties, Gil-Rodriquez & Fujii in Cassono & al. (2009:95)

[*Chondrophycus thyoides* (Kützing) Furnari (Furnari & al. 2001), *C. paniculatus* Furnari *Laurencia paniculata* (C. Agardh) J. Agardh]

ERYTHROCYSTIS J. Agardh

E. montagnei (Derbes & Solier) Silva

LAURENCIA Lamouroux

L. caduciramulosa Masuda & Kawaguchi

NOTE: Recorded from Rhodes and Saronikos (Tsiamis & al. 2011)

? *L. microcladia* Kützing

L. obtusa (Hudson) Lamouroux

L. pyramidalis Kützing

[*L. obtusa* (Hudson) Lamouroux var. *crucifera* Hauck]

OSMUNDEA Stackhouse

O. truncata (Kützing) Nam & Maggs (Nam, Maggs & Garbary 1994)
(*Laurencia* sp. sensu Athanasiadis 1987)

O. pelagosae (Schiffner) Nam

(*Rodriguezella pelagosae* Schiffner)

RODRIGUEZELLA Schmitz

R. ligulata *nomen*

NOTE: Kalogeroi islet (Tsiamis & al. 2010)

POLYSIPHONIEAE Schmitz & Falkenberg

ALSIDIUM C. Agardh

A. corallinum C. Agardh

A. helminthochortos (Schwendimann) Kützing

BOERGESENIELLA Kylin

Boergeseniella fruticulosa (Wulfen) Kylin
[P. fruticulosa (Wulfen) Sprengel]

DIGENEA C. Agardh

D. simplex (Wulfen) C. Agardh

HERPOSIPHONIA Nägeli

H. secunda (C. Agardh) Ambronnn

H. tenella (C. Agardh) Ambronnn

LOPHOCLADIA (J. Agardh) Schmitz

? L. lallemandii (Montagne) Schmitz

? L. trichoclados (C. Agardh) Schmitz

LOPHOSIPHONIA Falkenberg

L. cristata Falkenberg

L. obscura (C. Agardh) Falkenberg
[L. subadunca (Kützing) Falkenberg]

L. reptabunda (Suhr) Kylin

Lophosiphonia scopulorum (Harvey) Womersley
(P. scopulorum Harvey)

NEOSIPHONIA Kim & Lee

Neosiphonia sphaerocarpa (Børgesen) Kim & Lee.
[Polysiphonia sphaerocarpa Børgesen (Cormaci &al. 2004:171)]

OPHIOCLADUS Falkenberg
(Ophidocladus, *orth.var.*)

NOTE: Originally spelled *Ophidocladus*, but the first word is the masculine οφις (serpent, genitive οφε-ος), and hence the correct orthography should be *Ophiocladus*.

O. simpliciusculus (Croau *frat.* ex J. Agardh) Falkenberg

NOTE: Recorded from Andros Island (Tsiamis &al. 2010)

POLYSIPHONIA Greville

? P. biformis Zanardini

NOTE: *Dasya corallicola* Funk was reported from islands of the S. Archipelagus by Giaccone in 1968 (see Athanasiadis 1987). Following Feldmann, Funk (1961) considered this species a synonym of *Polysiphonia biformis* Zanardini

P. breviarticulata (C. Agardh) Zanardini

P. brodiei (Dillwyn) Sprengel

P. denudata (Dillwyn) Greville

P. elongata (Hudson) Sprengel

P. funebris J. Agardh

NOTE: Recorded from Rhodes and Saronikos Gulf (Tsiamis & al. 2011)

P. furcellata (C. Agardh) Harvey

P. opaca (C. Agardh) Moris & De Notaris

P. ornata J. Agardh

? P. sanguinea (C. Agardh) Zanardini

P. sertularioides (Grateloup) J. Agardh

P. subulifera (C. Agardh) Harvey

P. tenerima Kützing

P. cf. tripinnata J. Agardh

WOMERSLEYELLA Hollenberg

W. setacea (Hollenberg) R. Norris (Athansiadis 1997b)

POLYZONIEAE Schmitz

DIPTEROSIPHONIA Schmitz & Falkenberg

D. rigens (C. Agardh) Falkenberg

PTEROSIPHONIEAE Falkenberg

PTEROSIPHONIA Falkenberg

? P. complanata (Clemente) Falkenberg

? P. parasitica (Hudson) Falkenberg

? P. pennata (C. Agardh) Sauvageau

BONNEMAISONIALES J. Feldmann & Feldmann-Mazoyer

BONNEMAISONIACEAE Schmitz

ASPARAGOPSIS Montagne

? A. armata Harvey

[Falkenbergia cf. rufolanosa (Harvey) Schmitz]

NOTE: The record is only based on the sporophyte stage.

A. taxiformis (Delile) Trevisan

NOTE: First recorded in 1992 from Cape Drepanon in the North Aegean (Athansiadis 1997b: 475, as 'A. armata'; 1999a:250, as Asparagopsis gametophytes)

BONNEMAISONIA C. Agardh

B. asparagoides (Woodward) C. Agardh

B. clavata Hamel

GELIDIALES Kylin
GELIDIACEAE Kützing

BECKERELLA Kylin

B. mediterranea Huve

[*Ptilophora mediterranea* (Huve) R. Norris (1987)]

GELIDIUM Lamouroux

G. miniusculum (Weber-van Bosse) R. Norris

(*G. pusillum* var. *miniusculum* Weber van Bosse; Giaccone 1968)

G. pusillum (Stackhouse) Le Jolis

G. spinosum (Gmelin) Silva (Silva &al. 1996)

[*G. latifolium* (Greville) Bornet]

PTEROCLADIALLA Santelices & Hommersand

P. capillacea (Gmelin) Santelices & Hommersand (1997)

[*P. capillacea* (Gmelin) Bornet]

P. melanoidea (Bornet) Santelices & Hommersand (1997)

(*Gelidium melanoideum* Bornet)

GELIDIELLACEAE Fan

GELIDIELLA J. Feldmann & Hamel

G. antipae Celan

(*G. antipai*, orth. variant)

G. lubrica (Kützing) J. Feldmann & Hamel

G. nigrescens (J. Feldmann) J. Feldmann & Hamel (Dural &al. 1997)

G. tenuissima J. Feldmann & Hamel (Santelices & Rico 2002)

[*G. pannosa* (J. Feldmann) J. Feldmann & Hamel]

GIGARTINALES Schmitz
CALOSIPHONIACEAE Kylin

CALOSIPHONIA Crouan frat.

C. vermicularis (J. Agardh) Schmitz

SCHMITZIA Lagerheim

S. neapolitana (Berthold) Lagerheim

CAULACANTHACEAE Kützing

CAULACANTHUS Kützing

C. ustulatus (Turner) Kützing

CRUORIACEAE (J. Agardh) Kylin

CRUORIA Fries

C. cruriaeformis (Crouan frat.) Denizot

CYSTOCLONIACEAE Kützing

RHODOPHYLLIS Kützing

R. divaricata (Stackhouse) Papenfuss

DUDRESNAYACEAE J. Agardh

ACROSYMPHYTON Sjoestedt

A. purpuriferum (J. Agardh) Sjoestedt

DUDRESNAYA Crouan frat.

D. verticillata (Withering) Le Jolis

FURCELLARIACEAE Greville
FURCELLARIEAE Fries

HALARACHNION Kützing

H. ligulatum (Woodward) Kützing

NEUROCAULON Kützing

N. foliosum (Meneghini) Kützing

GIGARTINACEAE Bory

CHONDRACTHUS Kützing (Hommersand &al. 1993)

? *C. acicularis* (Roth) Fredericq
[*Gigartina acicularis* (Roth) Lamouroux]

C. teebei (Mertens in Roth) Kützing
[*Gigartina teebei* (Mertens in Roth) Lamouroux]

GLOIOSIPHONIACEAE Schmitz

THURETELLA Schmitz

T. schousboei (Thuret) Schmitz

HYPNEACEAE J. Agardh

HYPNEA Lamouroux

Hypnea anastomosans Papenfuss, Lipkin & Silva in Lipkin & Silva (2002)
(*H. esperi* Bory)

NOTE: Recorded from Saronikos Gulf and Rhodes (Tsiamis & Verlaque 2011)

? *H. cervicornis* J. Agardh

NOTE: *Hypnea cervicornis* is currently considered to be a synonym of *H. spinella* (C. Agardh) Kützing

H. musciformis (Wulfen) Lamouroux

H. valentiae (Turner) Montagne

NOTE: The original record from Rhodes by Reinbold in 1898 has been questioned (Tsiamis & Verlaque 2011), and is currently referred to *H. cornuta* (not previously recorded in the Aegean). *Hypnea valentiae* was recently reported from Rhodes Port (just outside the Aegean) by Tsiamis & Verlaque (l.c.)

KALLYMENIACEAE (J. Agardh) Funk

KALLYMENIEAE J. Agardh

KALLYMENIA J. Agardh

K. feldmanii Rodriguez-Prieto & Hommersand (2009)

NOTE: Collected in Sithonia by A. Manghisi (unpublished data)

K. lacerata J. Feldmann

? *K. reniformis* (Turner) J. Agardh

K. requienii (J. Agardh) J. Agardh (Cirik 1991)

MEREDITHIA J. Agardh

M. microphylla (J. Agardh) J. Agardh

(*Kallymenia microphylla* J. Agardh)

PEYSSONNELIACEAE Denizot

CRUORIELLA Crouan frat.

C. armorica Crouan frat.

HAEMATOCELIS J. Agardh

H. sp.

METAPEYSSONELIA Boudouresque, Coppejans & Marcot

M. feldmannii Boudouresque, Coppejans & Marcot

PEYSSONNELIA Decaisne

P. bornetii Boudouresque & Denizot

P. coriacea J. Feldmann

NOTE: Recorded from Rhodes port just outside the proper Aegean Sea (Tsiamis & al. 2010)

P. crispata Boudouresque & Denizot

P. dubyi Crouan frat.

P. harveyana J. Agardh

P. immersa Maggs & Irvine

P. polymorpha (Zanardini) Schmitz

P. rosa-marina Boudouresque & Denizot

P. rubra (Greville) J. Agardh

P. stoechas Boudouresque & Denizot

NOTE: Recorded from Karpathos (Catra & Giardina 2009) and Korinthiakos Gulf (Tsiamis &al. 2010); both sites neighbouring the Aegean

P. squamaria (Gmelin) Decaisne

P. sp.

(*P. codana* sensu Verlaque)

POLYSTRATA Heydrich

P. compacta (Foslie) Denizot ex Athanasiadis (1987)

NOTE: In introducing the new combination, Denizot did not cite page number

PHYLLOPHORACEAE Nägeli

GYMNOGONGRUS Martius

G. griffithsiae (Turner) Martius

PHYLLOPHORA Greville

P. nervosa (DC) Greville

[*P. crispa* Auctorum]

SCHOTTERA Guiry & Hollenberg

S. nicaeensis (Duby) Guiry & Hollenberg

RHIZOPHYLLIDACEAE Schmitz

CONTARINIA Zanardini

C. peyssonneliaeformis Zanardini

RHIZOPHYLLIS Kützing

R. squamariae (Meneghini) Kützing

SPHAEROCOCCACEAE Dumortier SPHAEROCOCCEAE J. Agardh

SPHAEROCOCCUS Stackhouse

S. coronopifolius Stackhouse

WURDEMANNIACEAE Taylor

WURDEMANNIA Harvey

W. miniata (Sprengel) J. Feldmann & Hamel

NEMASTOMATALES Kylin
 NEMASTOMATACEAE Schmitz
 NEMASTOMEAE J. Agardh

NEMASTOMA J. Agardh

N. dichotomum J. Agardh (Athanasiadis 1988)

SCHIZYMEMIACEAE Masuda & Guiry

PLATOMA (J. Agardh) Schmitz

P. cyclocolpum (Montagne) Schmitz
 (*P. cyclocarpa*, orth. variant; Athanasiadis 2000)

PREDAEA De Toni

P. ollivieri J. Feldmann (Athanasiadis 1988)

SCHIZYMEMIA J. Agardh

S. dubyi (Duby) J. Agardh

PLOCAMIALES (Kützing) Saunders & Kraft
 PLOCAMIACEAE Kützing

PLOCAMIUM Lamouroux

P. cartilagineum (Linné) Dixon

CRYPTONEMIALES Schmitz
 (HALYMEMIALES Saunders & Kraft)
 HALYMEMIACEAE Bory

ACRODISCUS Zanardini

A. vidovichii (Meneghini) Zanardini

CRYPTONEMIA J. Agardh

C. lomation (Bertoloni) J. Agardh

? *C. tunaiformis* (Bertoloni) Zanardini

GRATELOUPIA C. Agardh

G. filicina (Lamouroux) C. Agardh

HALYMEMIA C. Agardh

H. elongata C. Agardh

H. floresii (Clemente) C. Agardh
 (*H. floresia*, orth. variant)

H. latifolia Kützing

SEBDENIACEAE Kylin

SEBDENIA Berthold

S. rodrigueziana (J. Feldmann) Codomier ex Athanasiadis (1987) (Manghisi & Ribera 2007)

GRACILARIALES Fredericq & Hommersand
GRACILARIACEAE Nägeli

GRACILARIA Greville

G. bursa-pastoris (Gmelin) Silva

G. corallicola Zanardini

G. dura (C. Agardh) J. Agardh

? *G. gracilis* (Stackhouse) Steentoft, Irvine & Farnham (1995)
(*G. verrucosa* Auctorum p.p.)

RHODYMENIALES Schmitz
CHAMPIACEAE Kützing

CHAMPIA Desvaux

C. parvula (C. Agardh) Harvey

CHYLOCLADIA Greville

C. verticillata (Lightfoot) Bliding

GASTROCLONIUM Kützing

G. reflexum (Chauvin) Kützing

LOMENTARIACEAE J. Agardh

LOMENTARIA Lyngbye

? *L. articulata* (Hudson) Lyngbye

L. chylocladiella Funk

L. clavellosa (Turner) Gaillon

L. compressa (Kützing) Kylin

L. phalligera J. Agardh 1842
[*L. linearis* (Zanardini) Zanardini ex Kützing 1849]

RHODYMENIACEAE Harvey

BOTRYOCLADIA (J. Agardh) Kylin

B. boergesenii J. Feldmann

B. botryoides (Wulfen) J. Feldmann

CHRYSYMENIA J. Agardh

C. ventricosa (Lamouroux) J. Agardh

? FAUCHEA Bory & Montagne

? F. repens (C. Agardh) Montagne & Bory

RHODYMENIA Greville

R. sp. (Athanasiadis 1987)

NOTE: Ardissoni (Phycologia Mediterranea: 213. 1883) reported *R. palmetta* (Stackhouse) Greville from Smyrna.

MELANOPHYCEAE Rabenhorst
 (PHAEOPHYCEAE Kjellman)
 FUCOPHYCIDEAE Areschoug

DISCOSPORANGIALES O.C. Schmidt
 DISCOSPORANGIACEAE O.C. Schmidt

DISCOSPORANGIUM Falkenberg

Discosporangium mesarthrocarpum (Meneghini) Hauck (Kawai & al. 2007)

CUTLERIALES Kjellman
 CUTLERİACEAE Hauck

CUTLERIA Greville

C. adspersa (Mertens in Roth) De Notaris

C. chilosa (Falkenberg) Silva
 (C. monoica Ollivier; Dural & al. 1997)

C. multifida (Turner) Greville
 [Aglaozonia parvula (Greville) Zanardini]

ZANARDINIA Nardo ex Zanardini

Z. typus (Nardo) Silva in Greuter (2000)

FUCALES C. Agardh
 CYSTOSEIRACEAE De Toni
 CYSTOSEIROIDEAE Kjellman

CYSTOSEIRA C. Agardh

C. amentacea (C. Agardh) Bory

C. barbata (Stackhouse) C. Agardh

C. compressa (Esper) Gerloff & Nizamuddin

C. corniculata (Turner) Zanardini

C. crinita Duby

C. crinitophylla Ercegovic

C. dubia Valiante

C. elegans Sauvageau

C. foeniculacea (Linné) Greville
(*C. schiffneri* Hamel; Gomez Garreta 2001:133)

C. mediterranea Sauvageau

C. sauvageauana Hamel

C. spinosa Sauvageau
(*C. adriatica* Sauvageau; Gomez Garreta 2001:155)

C. susanensis Nizamuddin

NOTE: Recorded from Kalogeroi Islet (Tsiamis & al. 2010)

C. squarrosa De Notaris

C. zosteroides C. Agardh

SARGASSACEAE Kützing

SARGASSUM C. Agardh

S. acinarium (Linné) Setchell

S. hornschuchii C. Agardh

S. trichocarpum J. Agardh

S. vulgare C. Agardh

DESMARESTIALES Setchell & Gardner ARTHROCLADIACEAE Chauvin

ARTHROCLADIA Duby

A. villosa (Hudson) Duby

DICTYOTALES Hauck

DICTYOTACEAE Dumortier

DICTYOPTERIS Lamouroux

D. lucida Ribera Siguan, Gomez Garreta, Perez Ruzafa, Barcelo Marti & Rull Lluch
NOTE: Recorded from Rhodes port and Ladiko bay, just outside the proper Aegean Sea
(Tsiamis & al. 2010) and Smyrna (Taskin 2013).

D. polypodioides (D.C.) Lamouroux

DICTYOTA Lamouroux

Dictyota cyanoloma Tronholm, De Clerck, Gömez Garreta & Rull Lluch in Tronholm & al.
(2010).

NOTE: Recorded from Smyrna (Taskin 2013).

D. dichotoma (Hudson) Lamouroux

D. dichotoma var. intricata (C. Agardh) Greville
[D. implexa (Desfontaines) Lamouroux]

D. linearis (C. Agardh) Greville

DILOPHUS J. Agardh

D. fasciola (Roth) Howe
(D. mediterraneus Schiffner; De Clerck 2003:180)

D. fasciola var. repens (J. Agardh) J. Feldmann

D. spiralis (Montagne) Hamel

PADINACEAE Nägeli

PADINA Adanson

P. di-tristromatica N.-N. Win & H. Kawai in Win & al. (2011)

NOTE: Recorded from several localities in the Mediterranean including Lesbos. Molecular analyses cluster this species, and P. pavonicoidea N.-N. Win & H. Kawai, together with P. pavonica, but neither the variation of P. pavonica populations (in the Mediterranean-Caribbean) nor the identity of older names (presumed to be synonyms of P. pavonica) are clarified

P. pavonica (Linné) Lamouroux

(P. pavonia, orth. variant, P. mediterranea Bory)

P. pavonicoidea N.-N. Win & Kawai in Win & al. (2011)

NOTE: Recorded from Smyrna (Taskin 2013).

? P. tenuis Bory

NOTE: Also known as Padina boryana Thivy 1966, following Silva & al. (1966: 602) who interpreted P. tenuis Bory 1827 as a new combination [P. tenuis (C. Agardh) Bory] - the latter based on Bory's statement "...c'est le Zonaria Pavonia δ d'Agardh..". Womersley & Bailey (1970: 292-3) interpreted P. tenuis as a new species typified with the Commerson collection from Mauritius which is in agreement with Bory's intent to create "..le Padina tenuis, N., recueilli d'abord par Commerson vers l'embouchure de la grande rivière à l'Ile-de-France,..". Recognition of P. tenuis Bory precludes the possibility of changing the name of the species (in favor of Vaughaniella rupicola Børgesen 1950 - cited as a possible synonym by Silva & al. l.c.). Padina tenuis has been reported from several localities in the Aegean by Nizamuddin, but according to Win & al. (2011: 338) these records may be representative of Padina boergesenii Allender & Kraft.

STYPOPODIUM Kützing

S. schimperi (Kützing) Verlaque & Boudouresque (1991:201)

TAONIA J. Agardh

T. atomaria (Woodward) J. Agardh

ZONARIA C. Agardh

Z. tournefortii (Lamouroux) Montagne

ECTOCARPALES Oltmanns
 (CHORDARIALES Setchell & Gardner, DICTYOSIPHONALES Setchell & Gardner,
 SCYTOSIPHONALES J. Feldmann)

ACINETOSPORACEAE J. Feldmann

ACINETOSPORA Bornet

A. crinita (Carmichael) Kornmann

FELDMANNIA Hamel

F. battersioides (Ercegovic) Cormaci & Furnari (Dural & al. 1997)

F. globifera (Kützing) Hamel

F. irregularis (Kützing) Hamel

F. lebelii (Crohan frat.) Hamel

[*F. caespitula* var. *lebelii* (Crohan frat.) Knoepffler-Peguy (Taskin & Oezturk 2007:174, Ayvalik near Lesbos)]

F. padinae (Buffham) Hamel (Taskin & Oezturk 2007:175, Ayvalik near Lesbos)

F. paradoxa (Montagne) Hamel

HERPONEMA J. Agardh

H. graniferum Kuckuck

HINCKSIA J. Gray

(Giffordia Batters)

H. granulosa (Smith & Sowerby) Silva (Taskin & Oezturk 2007:175, Ayvalik near Lesbos)

H. mitchelliae (Harvey) Silva (Silva & al. 1987:73, 130)

[Giffordia mitchelliae (Harvey) Batters]

PYLAIELLA Bory

? *P. littoralis* (Linné) Kjellman

NOTE: Reported from the Makedonian coast by Anagnostidis (1968) with reservations. Older Mediterranean records by Hauck refer to *f. fluviatilis* that is apparently referable to Ectocarpus

SPONGONEMA Kützing

? *S. tomentosum* (Hudson) Kützing

NOTE: A report from the Pagasitikos Gulf [Diannelidis 1935, as *E. tomentosus* (Huds.) Lyngb.] is doubtful because 'Das Vorkommen liegt ausserhalb des Verbreitungsgebietes' (Gerloff & Geissler 1974:744)

CHORDARIACEAE Greville

CLADOSIPHON Kützing

C. irregularis (Sauvageau) Kylin (Taskin & Oezturk 2007:182, Ayvalik near Lesbos)

C. mediterraneus Kützing

GONTRANIA Sauvageau

G. lubrica Sauvageau

LIEBMANNIA J. Agardh

L. leveillei J. Agardh

MESOGLOIA C. Agardh

M. lanosa Crouan frat.

M. vermiculata (Smith & Sowerby) S.F. Gray (Taskin & Oezturk 2007:184, Ayvalik near Lesbos)

PHAEOSTROMA Kuckuck

P. bertholdii Kuckuck (Taskin & Oezturk 2007:178, Ayvalik near Lesbos)

SAUVAGEAUGLOIA Kylin

S. griffithsiana (Harvey) Hamel (Taskin & Oezturk 2007:184, Ayvalik near Lesbos)

ECTOCARPACEAE C. Agardh

ECTOCARPUS Lyngbye

E. siliculosus (Dillwyn) Lyngbye

E. siliculosus f. arctus (Kützing) Kuckuck

KUETZINGIELLA Kuckuck

K. battersii (Sauvageau) Kuckuck (Taskin & Oezturk 2007:178, Assos and Ayvalik near Lesbos)

ELACHISTACEAE Kjellman

ELACHISTA Duby

E. fucicola (Velle) Areschoug (Taskin & Oezturk 2007:185, Ayvalik near Lesbos)

E. intermedia Crouan frat.

HALOTHRIX Reinke

H. lumbicalis (Kützing) Reinke (Taskin & Oezturk 2007:185, Ayvalik near Lesbos)

GIRAUDIACEAE J. Feldmann

GIRAUDIA Derbes & Solier

G. sphacelarioides Derbes & Solier in Castagne

LEATHESIACEAE Farlow

CORYNOPHLOEA Kützing

NOTE: After κορυνη (head, club) and the masculine substantive φλοιος (cortex). Hence the

orthography should be *Corynophloea* (De Toni Sylloge Algarum vol.III. Fucoideae: 420. 1897). Translated to 'Keulenrinde' by Kützing in *Phycologia germanica*

C. flaccida (C. Agardh) Kützing

LEATHESIA S.F. Gray

? *L. mucosa* J. Feldmann

MICROCORYNE Strömfelt

? *M. ocellata* Strömfelt (Taskin 2008:176, Ayvalik near Lesbos)

MYRIACTULA Kuntze

M. rigida (Sauvageau) Hamel

M. rivulariae (Suhr) J. Feldmann

M. stellulata (Harvey) Levring

M. sp.

LITHODERMATACEAE Hauck

PSEUDOLITHODERMA Svedelius

P. adriaticum (Hauck) Verlaque

MYRIONEMATACEAE Nägeli

MYRIONEMA Greville

M. liechtensternii Hauck

M. orbiculare J. Agardh

M. strangulans Greville

M. sp.

MYRIOTRICHIACEAE Kjellman

MYRIOTRICHIA Harvey

M. adriatica Hauck

M. clavaeformis Harvey

[? *Streblonema sphaericum* (Derbes & Solier) Thuret]

M. repens Hauck

PROTASPEROCOCCUS Sauvageau

P. myriotrichiformis Sauvageau

PETROSPONGIACEAE Racault & al.

CYLINDROCARPUS Crouan frat.

C. kuckuckii Taskin &al. (2010: 264, Ayvalik near Lesbos)

PUNCTARIACEAE (Thuret) Kjellman

ASPEROCOCCUS Lamouroux

A. bullosus Lamouroux

A. ensiformis (Delle Chiaje) Wynne (Taskin 2008:179, Ayvalik near Lesbos)

A. fistulosus (Hudson) Hooker

PUNCTARIA Greville

P. latifolia Greville

SCYTOSIPHONACEAE Farlow

COLPOMENIA (Endlicher) Derbes & Solier

C. peregrina Sauvageau (Orfanidis 1990)

C. sinuosa (Mertens in Roth) Derbes & Solier

HYDROCLATHRUS Bory

H. clathratus (C. Agardh) Howe

PETALONIA Derbes & Solier

P. fascia (O. F. Mueller) Kuntze

NOTE: The sporophyte stage is considered identical with *Ralfsia clavata* (Harvey) Crouan frat.

SCYTOSIPHON C. Agardh

S. lomentaria (Lyngbye) Link

NOTE: The sporophyte stage is considered identical with *Microspongium gelatinosum* Reinke

SOROCARPACEAE Pedersen

KUCKUCKIA Hamel

K. spinosa (Kützing) Kuckuck

SPERMATOCHNACEAE Kjellman

NEMACYSTUS Derbes & Solier

N. flexuosus var. *giraudyi* (J. Agardh) de Jong in de Jong & Prud'homme van Reine (1997:15)

SPERMATOCHNUS Kützing

S. paradoxus (Roth) Kützing

STILOPHORACHEAE Nägeli

STILOPHORA J. Agardh

S. tenella (Esper) Silva (Silva, Basson & Moe 1996)
[*S. rhizodes* (Turner) J. Agardh]

STRIARIACEAE Kjellman

STICTYOSIPHON Kützing

S. ardiaticus Kützing (Taskin 2008:182, Ayvalik near Lesbos)

STRIARIA Greville

S. attenuata (Greville) Greville

LAMINARIALES Kylin
PHYLLARIACEAE Tilden
PHYLLARIOIDEAE Kjellman

SACCORHIZA Pylaie

S. polyschides (Lightfoot) Batters

NOTE: Only known from a single record from Chios

RALFSIALES Lim & Kawai
RALFSIACEAE Farlow

RALFSIA Berkeley

R. verrucosa (Areschoug) Areschoug

SPOROCHNALES Sauvageau
SPOROCHNACEAE Greville

NEREIA Zanardini

N. filiformis (J. Agardh) Zanardini

SPOROCHNUS C. Agardh

S. pedunculatus (Hudson) C. Agardh

SPHACELARIALES Migula
CHORISTOCARPACEAE Kjellman

CHORISTOCARPUS Zanardini

C. tenellus (Kützing) Zanardini

CLADOSTEPHACEAE Oltmanns

CLADOSTEPHUS C. Agardh

NOTE: After κλαδος (branch) and the neuter substantive στεφος (Lyngbye in Tentamen Hydrophytologiae Danicae:102. 1819)

C. spongiosum f. *verticillatum* (Lightfoot) Prud'homme van Reine

DISCOSPORANGIACEAE O.C. Schmidt

DISCOSPORANGIUM Hauck

D. mesarthrocarpum (Meneghini) Hauck (Dural & al. 1997)

SPHACELARIACEAE Decaisne

SPHACELARIA Lyngbye

S. cirrosa (Roth) C. Agardh

S. fusca (Hudson) S.F. Gray

S. plumula Zanardini

S. rigidula Kützing (Dural & al. 1997)

S. tribuloides Meneghini

STYPOCAULACEAE Oltmann

HALOPTERIS Kützing

H. filicina (Grateloup) Kützing

STYPOCAULON Kützing

S. scoparium (Linné) Kützing

CHLOROPHYCEAE Kützing
TETRASPORALES Lemmermann
TETRASPORACEAE (Nägeli) Wittrock

PALMOPHYLLUM Kützing

P. crassum (Naccari) Rabenhorst

ULOTRICHALES Borzi
ULOTRICHACEAE Kützing

ULOTHRIX Kützing

? U. flacca (Dillwyn) Thuret
(U. pseudoflacca Wille)

U. implexa (Kützing) Kützing

? U. subflaccida Wille (Dural & al. 1997, Smyrna)

CTENOCLADALES P.C.Silva
ULVELACEAE Schmidle

ACROCHAETE Pringsheim

A. inflata (Ercegovic) Gallardo & al. (Panayotidis 1979)
(Pseudodictyon infatum Ercegovic)

ENDOCLADIA Reinke

E. leptochate (Huber) Burrows
[Acrochaete leptochate (Huber) Nielsen]

E. viridis Reinke
[Acrochaete viridis (Reinke) Nielsen, Endoderma viride (Reinke) Lagerheim]

EPICLADIA Reinke

E. sp.

PRINGSHEIMIELLA von Hoehnel

P. scutata (Reinke) Marchewianka

PHAEOPHILALES Chappell &al. (1990)
PHAEOPHILACEAE Chappell &al.

PHAEOPHILA Hauck

P. dendroides (Crouan frat.) Batters

ACROSIPHONIALES Jonsson
ACROSIPHONIACEAE Jonsson

UROSPORA Areschoug

U. penicilliformis (Roth) Areschoug
(Codiolum gregarium Braun)

SPONGOMORPA Kützing

? S. aeruginosa (Linné) Hoek
[Cladophora lanosa (Roth) Kützing, Cladophora lanosa f. uncialis (O. F. Mueller) Hauck]

? S. arcta (Dillwyn) Kützing
[Cladophora arcta (Dillwyn) Kützing]

ULVALES Blackman & Tansley
GOMONTIACEAE De Toni

GOMONTIA Bornet & Flahault

G. polyrhiza (Lagerheim) Bornet & Flahault

MONOSTROMATACEAE Kunieda

BLIDINGIA Kylin

B. marginata (J. Agardh) Dangeard
(Enteromorpha marginata J. Agardh)

ULVACEAE Dumortier

ENTEROMORPHA Link

E. clathrata (Roth) Greville

E. compressa (Linné) Nees

E. flexuosa (Wulfen) J. Agardh
 (*E. juergensii* Kützing, *E. lingulata* J. Agardh)

E. intestinalis (Linné) Nees

E. linza (Linné) J. Agardh

E. multiramosa Bliding 1963

NOTE: Bliding (*Opera Botanica* 8: 116. 1963) specified that his account was based on 'specimens from Naples' and in particular on a 'mother plant (Bliding 1960 fig.4a..)' that was cultured in Sweden. This statement fulfills the requirement of 'indication of type...an entire gathering..even if it consists of two or more specimens...' (Art.40.2 and 40.3. Note 2). The type of *E. multiramosa* is the illustrated material on a single herbarium sheet (Bliding in *Bot. Notiser.* 113: 117-9, f.4.a. 1960), collected in 'Naples: March 1958..in the harbour of S:a Lucia, growing on Cladophoras' (Bliding 1963: 116). This material was the start of Bliding's culture experiments that produced his account in 1963. A Latin diagnosis was provided via 'Ref.: Bliding 1960, p.177' as also the herbarium holding the type (Bliding 1963: 116). Hence *E. multiramosa* Bliding 1963 is a valid name. The species should be transferred to the 'clathrata-group' when that information has also been evaluated. The wide range of habitats and diminutive size of the species is in accord with its cosmopolitan distribution that includes Australia, N. Zealand, the Indian Ocean, Papua New Guinea, Fiji and Canary Islands, Brazil and even the Caribbean (Athanasiadis 1987: 140)

E. muscoides (Clemente) Cremades

[*E. ramulosa* (Smith & Sowerby) Carmichael, *E. crinita* Nees]

E. prolifera (O. F. Mueller) J. Agardh

ULVA Linné

U. fasciata Delile

U. lactuca Linné

U. lactuca f. *laciiniata* (J. Agardh) De Toni
 (*U. rigida* f. *laciiniata* J. Agardh)

U. rigida C. Agardh

[*U. lactuca* var. *rigida* (C. Agardh) Le Jolis]

U. rotundata Bliding

CLADOPHORALES G. S. West
 ANADYOMENACEAE Kützing

ANADYOMENE Lamouroux

A. stellata (Wulfen) C. Agardh
 (*Ulva stellata* Wulfen in Jacquin, *A. flabellata* Lamouroux)

MICRODICTYON Decaisne

M. tenuius J. Gray
 [*M. umbilicatum* (Vellley) Zanardini p.p.]

CHAETOSIPHONIACEAE Blackman & Tansley

BLASTOPHYSA Reinke

B. rhizopus Reinke
(B. polymorpha Kjellman)

CLADOPHORACEAE Willie

CHAETOMORPHA Kützing

C. aerea (Dillwyn) Kützing

C. crassa (C. Agardh) Kützing

C. linum (O.F. Mueller) Kützing
[Conferva linum O.F.Mueller, C. chlorotica (Montagne) Kützing]

CLADOPHORA Kützing

C. albida (Nees) Kützing

[C. hamosa (Kützing) Kützing, C. neesiorum (C. Agardh) Kützing, C. pumila Kützing]

C. coelothrix Kützing

[C. repens (J. Agardh) Harvey, Cladophoropsis modonensis (Kützing) Reinbold,
Siphonocladus concrescens Reinbold (Leliart & Coppejans 2006:672)]

C. dalmatica Kützing

(C. arachnoidea Schiffner)

C. echinus (Biasoletto) Kützing

[C. cornea (Kützing) Kützing]

C. feredayi Harvey

(C. pellucida f. nana Hauck)

C. flexuosa (O. F. Mueller) Kützing

(C. gracilis Kützing)

C. glomerata (Linné) Kützing

C. hutchinsiae (Dillwyn) Kützing

(C. rissoana Kützing)

C. laetevirens (Dillwyn) Kützing

[C. meneghiniana (Kützing) Kützing, C. repens Harvey f. meneghiniana (Kützing) Hauck, C.
utriculosa var. laetevirens (Dillwyn) Hauck, ? C. affinis Schiffner, Siphonocladus rhodensis
Reinbold (Leliart & Coppejans 2006:673)]

C. lehmanniana (Lindenberg) Kützing

[C. utriculosa Kützing, C. utriculosa var. longiarticulata Kützing, C. utriculosa var. ramulosa
(Meneghini) Hauck, C. ramulosa Meneghini]

C. nigrescens Zanardini ex Frauenfeld

NOTE: Recorded from Andros (Tsiamis &al. 2010)

C. pellucida (Hudson) Kützing

[C. trichotoma (C. Agardh) Kützing]

C. prolifera (Roth) Kützing
[*C. prolifera* var. *scoparia* (Kützing) Schiffner]

C. retroflexa (Crouan frat.) van den Hoek

C. rupestris (Linné) Kützing
[*C. rupestris* var. *mediterranea* Kützing, *C. rupestris* f. *mediterranea* (Kützing) Ardisson]

C. sericea (Hudson) Kützing
[*C. rudolphiana* (C. Agardh) Harvey, *C. mediterranea* Hauck]

? *C. socialis* Kützing

C. vadorum (Areschoug) Kützing
(*C. corynartha* Kützing)

C. vagabunda (Linné) Hoek
[*C. fracta* f. *marina* Hauck, *C. penicillata* Kützing, *C. utriculosa* var. *aegaea* (Kützing)
Kützing]

RHIZOCLONIUM Kützing

R. riparium (Roth) Harvey

R. tortuosum (Dillwyn) Kützing
[*Chaetomorpha tortuosa* Kützing, *Chaetomorpha capillaris* (Kützing) Børgesen]

SIPHONOCLEDACEAE Schmitz

CLADOPHOROPSIS Børgesen

? *C. membranacea* (C. Agardh) Børgesen
[? *Siphonocladus membranaceus* (C. Agardh) Bornet var. *caespitosus* (C. Agardh) De Toni, ?
Siphonocladus psyttaliensis Schmitz]

Cladophoropsis sundanensis Reinbold (Leliart & Coppejans 2006:679)

SIPHONOCLEDUS Schmitz

S. pusillus (Kützing) Hauck
(*Valonia pusilla* Kützing, *S. wilbergii* Schmitz)

VENTRICARIA Olsen & J. West

V. ventricosa (J. Agardh) Olsen & West
(*Valonia ventricosa* J. Agardh)

VALONIACEAE Kützing

VALONIA C. Agardh

V. macrophysa Kützing

Valonia utricularis (Roth) C. Agardh
(*Conferva utricularis* Roth, ? *V. caespitula* Zanardini)

BRYOPSIDALES Schaffner BRYOPSIDACEAE Bory

BRYOPSIS Lamouroux

B. adriatica (J. Agardh) Meneghini
[*B. plumosa* (Hudson) C. Agardh var. *adriatica* (J. Agardh) Hauck]

B. duplex De Notaris
[*B. balbisiana* Lamouroux sensu J. Feldmann, *B. balbisiana* Lamouroux var. *disticha* J. Agardh, *B. disticha* (J. Agardh) Kützing]

B. corymbosa J. Agardh

B. feldmannii Gallardo & Furnari
(*B. cupressoides* J. Feldmann)

B. hypnoides Lamouroux

B. muscosa Lamouroux

B. plumosa (Hudson) C. Agardh

B. sp.

CAULERPACEAE Kützing

CAULERPA Lamouroux

C. prolifera Lamouroux

NOTE: Recorded from Crete

C. cylindracea Sonder

[*C. racemosa* (Forsskål) J. Agardh var. *cylindracea* (Sonder) Verlaque, Huisman & Boudouresque (2003:336); *C. racemosa* cf. var. *occidentalis* (J. Agardh) Børgesen (Verlaque &al. 2000, tab. II; Panayotidis & Zuljevic 2001)]

NOTE: Recorded from Crete, Kalimnos, Samos, Saronikos, Tristinika (Sithonia, June 2009; Tsiamis &al. 2014)

C. racemosa var. *lamourouxii* f. *requienii* (Montagne) Weber van Bosse (Verlaque &al. 2000, table II)

NOTE: Recorded from the Island of Thira

CODIACEAE Kützing

CODIUM Stackhouse

? *C. adhaerens* C. Agardh

C. bursa (Olivi) C. Agardh

C. coralloides (Kützing) Silva

C. decorticatum (Woodward) Howe

[*C. elongatum* (Turner) C. Agardh, *Fucus elongatus* Turner]

C. effusum (Rafinesque) Delle Chiaje

(*C. difforme* Kützing)

C. fragile subsp. *tomentosoides* (Goor) Silva

C. vermilara (Olivi) Delle Chiaje

DERBECIACEAE Hauck

DERBESIA Solier

D. cf. tenuissima (Morris & De Notaris) Crouan frat.
(*Halicystis* cf. *parvula* Schmitz)

PEDOBESIA MacRaild & Womersley

P. simplex (Kützing) Wynne & Leliaert (2001)
[P. lamourouxii (J. Agardh) J. Feldmann & al., *Derbesia lamourouxii* (J. Agardh) Solier, ?
Derbesia cervicornis Schiffner]

HALIMEDACEAE Link

HALIMEDA Lamouroux

H. tuna (Ellis & Solander) Lamouroux

PENICILLUS Lamarck

P. capitatus Lamarck
(*Espera mediterranea* Decaisne)

UDOTEACEAE J. Agardh

FLABELLIA Reichenbach

F. petiolata (Turra) Nizamuddin
[*Udotea petiolata* (Turra) Børgesen, *Udotea desfontainii* (Lamouroux) Decaisne,
? *Udotea minima* Ernst]

PSEUDOCHLORODESMIS Børgesen

P. furcellata (Zanardini) Børgesen
(*Bryopsis furcellata* Zanardini)

DASYCLADALES Pascher
DASYCLADACEAE Kützing

DASYCLADUS C. Agardh

D. vermicularis (Scopoli) Krasser
[D. clavaeformis (Roth) C. Agardh, *Myrsidrum bertolonii* Bory]

ACETABULARIACEAE Nägeli

ACETABULARIA Lamouroux

A. acetabulum (Linné) Silva
(A. mediterranea Lamouroux)

POLYPHYSCACEAE Kützing

PARVOCAULIS Berger, Fettweiss, Gleissberg, Liddle, Richter, Sawitzky & Zuccarello
(2003)

P. parvula (Solms-Laubach) Berger & al. (2003)
 [Polyphysa parvula (Solms-Laubach) Schnetter & Bula-Meyer, Acetabularia parvula Solms-Laubach (Cirik 1978), Acetabularia moebii Solms-Laubach, ? Acetabularia wettsteinii Schussnig]

REFERENCES

- Athanasiadis A.** 1987. A survey of the seaweeds of the Aegean Sea with taxonomic studies on species of the tribe Antithamnieae (Rhodophyta).- Goterna, Kungaev. 174 pp
- 1988. North Aegean Marine Algae II. Studies on the thallus structure and reproduction of *Nemastoma dichotomum* J. Agardh and *Predaea ollivieri* J. Feldmann (Rhodophyta, Gigartinales).- Bot. mar. 31:23-32
 - 1989. North Aegean Marine Algae III. Structure and development of the encrusting coralline *Titanoderma cystoseirae* (Rhodophyta, Lithophylloideae).- Nord. J. Bot. 9:435-441
 - 1995. Morphology, anatomy and reproduction of the eastern Mediterranean coralline *Tenarea tortuosa* and its relationship to members of the Lithophylloideae and Mastophoroideae (Rhodophyta, Corallinales).- Nord. J. Bot. 15:655-663
 - 1996. Morphology and classification of the Ceramioideae (Rhodophyta) based on phylogenetic principles.- Opera Bot. 128:1-216
 - 1997a. On the typification and taxonomic status of *Melobesia notarisii* Dufour (Rhodophyta, Corallinales).- Phycologia 36:410-415
 - 1997b. North Aegean Marine Algae IV. *Womersleyella setacea* (Hollenberg) R. E. Norris (Rhodophyta, Ceramiales).- Bot. mar. 40:473-476
 - 1999a. *Mesophyllum macedonis* nov. sp. (Rhodophyta, Corallinales), a putative Tethyan relic in the North Aegean Sea.- Eur. J. Phycol. 34:239-252
 - 1999b. The taxonomic status of *Lithophyllum stictaeforme* (Rhodophyta, Corallinales) and its generic position in light of phylogenetic considerations.- Nord. J. Bot. 19:735-745
 - 2000. Proposal to conserve the name *Platoma* (Rhodophyta) as being of neuter gender.- Taxon 49:809-811
 - 2003. Taxonomy and systematics of Rhodophyta with reference to the Mediterranean taxa.- Fl. Medit. 12 [2002]:93-168
 - & Neto A.I. 2010. On the occurrence of *Mesophyllum expansum* (Philippi) Cabioch & Mendoza (Melobesioideae, Corallinales, Rhodophyta) in the Mediterranean, the Canary Isles and the Azores.- Bot. mar. 53:333-341
- Aysel V.** 1983. The Chondria Agardh species (Rhodophyta, Ceramiales) of the Aegean coast.- Doga Bilim Dergisi:Temel Bilm.:Cilt 7:47-57
- Babbini L. & Bressan G.** 1997. Recensement de Corallinacees de la Mer Mediteranee & considerations phytogeographiques.- Bibl. Phycol. 103:1-421
- Basso D.** 1995. Living calcareous algae by a paleontological approach: The genus *Lithothamnion* Heydrich nom. cons. from the soft bottoms of the Tyrrhenian Sea (Mediterranean).- Riv. Ital. Paleont. Strat. 101: 349-366.
- , - & Bressan G. 2010. A re-description of *Lithothamnion crispatum* and the status of *Lithothamnion superpositum* (Rhodophyta, Corallinales). Phycologia 50: 144-155.
- Berger S., Fettweiss U., Gleissberg S., Liddle L.B., Richter U., Sawitzky H. & Zuccarello G.S.** 2003. 18S rDNA phylogeny and evolution of cap development in Polyphysaceae (formerly Acetabulariaceae; Dasycladales, Chlorophyta).- Phycologia 42:506-561
- Brodie J., Bartsch I., Neefus C., Orfanidis S., Bray T. & Mathieson T.** 2007. New insights into the cryptic diversity of the North Atlantic- Mediterranean 'Porphyra leucosticta' complex: *P. olivii* sp.nov. and *P. rossengurttii* (Bangiales, Rhodophyta).- Eur.J.Phycol. 42:3-28
- Cassano V., Diaz-Larrea J., Senties A., Oliveira M.C., Candelaria Gil-Rodriguez M. & Fujii M.T.** 2009. Evidence for the conspecificity of *Palisada papillosa* with *P. perforata* (Ceramiales, Rhodophyta) from the western and eastern Atlantic Ocean on the basis of morphological and molecular analyses.- Phycologia 48:86-100
- Catra M. & Giardina S.** 2009. A survey of the marine macroalgae of Karpathos island (the Aegean Sea, Greece).- Plant Biosystems 143:509-515
- Chappell D.F., O'Kelly C.J., Wilcox L.W. & Floyd G.L.** 1990. Zoospore flagellar apparatus architecture and the taxonomic position of *Phaeophila dendroides* (Ulvophyceae, Chlorophyta).- Phycologia 29:515-523

- Cirik S.** 1978. Recherches sur la vegetation marine des cotes turques de la mer Egee. Etudes particulières des Peyssonneliacees de Turquie.-These de 3eme cycle, Universite Pierre & Marie Curie, Paris VI, France
- 1991. A propos de la vegetation marine de la baie d'Akkuyu (Mersin, Turquie).- Flora Mediterranea 1:205-212
- Cormaci M., Furnari G., Giaccone G. & Serio D.** 2004. Alien macrophytes in the Mediterranean Sea:A review.- Recent Res. Devel. Environ.Biol. 1:153-202.
- De Clerck O.** 2003. The genus Dictyota (Dictyotales, Phaeophyta) in the Indian Ocean.- Opera bot. Belgica 13:1-205
- De Jong Y.S.D.M.** 1997. Studies on Dasyaceae I. Reproductive structures of *Eupogodon planus* and *E. spinellus* (Ceramiales, Rhodophyta) from Corsica (Mediterranean).- Phycologia 36:281-292
- & **Prud'homme van Reine W.F.** 1997. A review of the genus *Nemacystus* (Spermatocladaceae, Chordariales, Phaeophyceae), including phylogenetic and biogeographical hypotheses.- Nova Hedwigia 64:1-40
- Dural B., Aysel V., Loek A. & Guner H.** 1997. Benthic algal flora of the natural and artificial substrata of Hekim Island (Izmir, Turkey).- Algol. Studies 85:31-48
- Giaccone G.** 1968. Raccolte di fitobenthos nel Mediterraneo orientale.- Giorn. bot. ital. 102:217-228
- Gomez Garreta A.** (ed.) 2001. Flora Phycologica Iberica Vol. 1. Fucales.- Universidad de Murcia
- Gerloff J. & Geissler U.** 1974. Eine revidierte Liste der Meeresalgen Griechenlands.- Nova Hedwigia 12:721-793
- Greuter W.** 2000. Notices of Publications, Cryptogamae:2.- OPTIMA Newsletter 35:1-36
- Funk G.** 1961. Ergänzungen und Berichtigungen zur Meeresalgenflora von Neapel.- Pubbl. Staz. Zool. Napoli 32:172-186
- Furnari G., Cormaci M. & Serio D.** 2001. The Laurencia complex (Rhodophyta, Rhodomelaceae) in the Mediterranean Sea:an overview.- Crypt.Algol. 22:331-373
- Hamel G.** 1931. Sur le Cladostephus dubius Bory.- I:Travaux Cryptogamique dedies a Louis Mangin, 309-312, Paris
- Hommersand M.H., Guiry M.D., Fredericq S. & Leister G.L.** 1993. New perspectives in the taxonomy of the Gigartinaceae (Gigartinales, Rhodophyta).- Hydrobiologia 260/261:105-120
- Huisman J.M. & Borowitzka M.A.** 1990. A revision of the Australian species of *Galaxaura* (Rhodophyta, Galaxauraceae), with a description of *Tricleocarpa* gen. nov.- Phycologia 29:150-172
- & **Townsend R.A.** 1993. An examination of Linnaean and pre-Linnaean taxa referable to *Galaxaura* and *Tricleocarpa* (Galaxauraceae, Rhodophyta).- Bot. J. Linnaean Soc. 133:95-101
- Irvine L.M. & Chamberlain Y.M.** 1994. Seaweeds of the British Isles. Volume 1 Rhodophyta, Part 2B Corallinales, Hildenbrandiales.- HMSO, London
- Kawai H. , Hanyuda T., Draisma S.G.A. & Mueller D.G.** 2007. Molecular phylogeny of *Discosporangium mesarthrocarpum* (Phaeophyceae) with a reinstatement of the order Discosporangiales.- J.Phycol. 43:186-194
- Lazaridou E. & Budouresque C.-F.** 1992. On a new species of Ceramium, *C. graecum* (Rhodophyta:Cramiaceae) from Greece.- Bot. mar. 35:561-565
- Leliaert F. & Coppejans E.** 2006. A revision of *Cladophoropsis* Børgesen (Siphonocladales, Chlorophyta).- Phycologia 45:657-679
- Lemoine M.** 1973. Contribution a etude du genre *Lithoporella* (Corallinacees).- Rev. Algol. 11:42-57
- Lipkin Y. & Silva P.S.** 2002. Marine algae and seagrasses of the Dahlak Archipelago, southern Red Sea.- Nova Hedwigia 75:1-90
- Maggs C.A. & Hommersand M.H.** 1993. Seaweeds of the British Isles. Vol. 1 Rhodophyta, Part 3 A Ceramiales.- HMSO, London
- Manghisi A. & Ribera M.A.** 2007. Lectotypification of *Halymenia rodrigueziana* J. Feldmann (= *Sebdenia rodrigueziana* (J. Feldmann) Codomier ex Athanasiadis (Sebdeniaceae, Rhodophyta)).- Ann. J. Bot. Madrid 64:75-78
- Nam K.W.** 2007. Validation of the generic name *Palisada* (Rhodomelaceae, Rhodophyta).- Algae 22:53-55
- , **Maggs C.A. & Garbary D.J.** 1994. Resurrection of the genus *Osmundea* with an emendation of the generic delineation of *Laurencia* (Ceramiales, Rhodophyta).- Phycologia 33:384-395

- Norris R.E.** 1987. A re-evaluation of Ptilophora Kützing and Beckerella Kylin (Gelidiales, Rhodophyceae) with a review of South African Species.- Bot. mar. 30:243-258
- 1993. Taxonomic studies on Ceramiaeae (Ceramiales, Rhodophyta) with predominantly basipetal growth of corticating filaments.- Bot. mar. 36:389-398
- Orfanidis S.** 1990. Temperature and photoperiodic responses of several Mediterranean macroalgae in relation to their distribution in the North Atlantic Ocean. - PhD, Aristotle Univ. of Thessaloniki (in Greek with English summary)
- Panayotidis P.** 1979. Etude phytosociologique de deux aspects saisonniers de la flore epiphyte des feuilles de *Posidonia oceanica* (Linnaeus) Delile, dans le golfe de Thessaloniki (Mer Egee, Grece).- Thalassographica 3:93-104
- & **Zuljevic A.** 2001. Sexual reproduction of the invasive green alga *Caulerpa racemosa* var. *occidentalis* in the Mediterranean Sea.- Oceanol. Acta 24:199-203
- Santelices B. & Hommersand M.H.** 1997. *Pterocladiella*, a new genus in the Gelidiaceae (Gelidiales, Rhodophyta).- Phycologia 36:114-119
- & **Rico J.M.** 2002. Nomenclature and typification of *Gelidiella tenuissima* (Gelidiales, Rhodophyta).- Phycologia 41:436-440
- Sartoni G. & Boddi S.** 2002. *Ceramium bisporum* (Ceramiaceae, Rhodophyta), a new record for the Mediterranean algal flora.- Bot. mar. 45:566-570
- & **De Biasi A.M.** 1999. A survey of the marine algae of Milos Island, Greece.- Crypt. Algol. 20:271-283
- Serio D., Cormaci M., Furnari G. & Boisset F.** 2010. First record of *Palisada maris-rubri* (Ceramiales, Rhodophyta) from the Mediterranean Sea along with three proposed transfers to the genus *Palisada*.- Phycological Research 85:9-16
- Silva P. C., Basson P.W. & Moe R.L.** 1996. Catalogue of the Benthic Marine Algae of the Indian Ocean.- Univ. Cal. Publ. Bot. 79:1-1259
- , **Menez E.G. & Moe R.L.** 1987. Catalog of the Benthic Marine Algae of the Philippines.- Smiths. Contr. Mar. Sci. 27:iv + 1-179
- Steentoft M., Irvine L.M. & Farnham W.F.** 1995. Two terete species of *Gracilaria* and *Gracilariaopsis* (Gracilariales, Rhodophyta) in Britain.- Phycologia 34:113-127
- Tronholm A., Steen F., Tyberghein L., Leliaert F., Verbruggen H., Ribera Siguán M.A. & De Clerck O.** 2010. Species delimitation, taxonomy, and biogeography of *Dictyota* in Europe (Dictyotales, Phaeophyceae).- J. Phycol. 46: 1301-1321
- Taskin E.** 2008. The marine brown algae of the East Aegean Sea and Dardanelles II. Ectocarpaceae, Chordariaceae, and Scytoniphonaceae.- Crypt., Algol. 29:173-186
- 2013. New records of three dictyotalean brown algae for Turkey.- Bot. mar. 56: 299-302.
- & **Oztürk M.** 2007. The marine brown algae of the East Aegean Sea and Dardanelles I. Ectocarpaceae, Pylaiellaceae, Chordariaceae, Elachistaceae and Giraudiaceae.- Crypt., Algol. 28:169-190
- **Wynne M. & Oztürk M.** 2010. *Cylindrocarpus kuckuckii* sp.nov. (Chordariaceae, Phaeophyceae), a newly recognized species from the Aegean Sea (Turkey).- Nova Hedwigia 90: 263-270
- Tsiamis K., Montesanto B., Panayotidis P. & Katsaros C.** 2011. Notes on new records of red algae (Ceramiales, Rhodophyta) from the Aegean Sea (Greece, eastern Mediterranean).- Plant Biosystematics 145: 873-884.
- , - , **Panayotidis P. & Montesanto B.** 2010. New macroalgal records for the Aegean Sea (Greece, eastern Mediterranean Sea).- Bot.marina 53:319-331
- , **Panayotidis P., Economou-Amilli A. & Katsaros C.** 2014. Seaweeds of the Greek coasts. II. Ulvophyceae.- Mediterranean Marine Science. 15: 000-000
- & **Verlaque M.** 2011. A new contribution to the alien red macroalgal flora of Greece (eastern Mediterranean) with emphasis on *Hypnea* species.- Crypt.Algal.32: 393-410
- Verlaque M. & Boudouresque C.-F.** 1991. *Styropodium schimperi* (Buchinger ex Kützing) Verlaque & Boudouresque comb. nov., algue de mer Rouge recemment apparue en Méditerranée.- Crypt., Algol. 12:195-211
- , **Meinesz A & Gravez V.** 2000. The *Caulerpa racemosa* complex (Caulerpales, Ulvophyceae) in the Mediterranean Sea.- Bot. mar. 43:49-68
- , **Durand C., Huisman J., Boudouresque C.-F. & Le Parco Y.** 2003. On the identity and origin of the Mediterranean invasive *Caulerpa racemosa* (Caulerpales, Chlorophyta).- Eur. J. Phycol. 38:325-339

- Win N.-N., Hanyuda T., Draisma S.G.A., Furnari G., Meinesz A. & Kawai H.** 2011. Padina ditriformatica sp. nov. and Padina pavonicoidea sp. nov. (Dictyotales, Phaeophyceae), two new species from the Mediterranean Sea based on morphological and molecular markers.- Eur. J. Phyc. 46:327-341
- Wynne M.J.** 1991. A change in the name of the type of Chondria C. Agardh (Rhodomelaceae, Rhodophyta).- Taxon 40:316-318
- & **Leliaert F.** 2001. Pedobesia simplex (Kützing) comb. nov. (Chlorophyta), a new name for P. lamourouxii and its first report from the Indian Ocean.- Crypt. Algol. 22:3-14

NOTE: This is a periodically updated electronic file, to be regarded as a preliminary version not effectively published (Art. 30.2).